



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

370. Proposed by R. C. ARCHIBALD, Paris, France.

The trisectors of the angles of any triangle  $ABC$  are, in order,  $AF, AE, CE, CD, BD, BF$ . Show synthetically that  $D, E, F$  are the vertices of an equilateral triangle.

Solution by A. H. HOLMES, Bruntwick, Me.



Let  $ABC$  be the triangle with  $AE, AF$  trisectors of  $\angle BAC$ ;  $BF, BD$  trisectors of  $\angle ABC$ , and  $CE, CD$  trisectors of  $\angle ACB$ . Draw  $DE, DF$ , and  $EF$ . Then  $DEF$  is an equilateral triangle. From  $D$  draw  $DG$  parallel to  $AF$  and cutting  $AE$  in  $G$ . Take, on  $AF$ ,  $AH = DG$ . Then  $\angle GDH = \angle EAF = \frac{1}{3} \angle BAC$ .

Suppose the triangle  $GED$  to be moved so that  $GE$  is colinear with  $AF$  and the point  $E$  is at the point  $F$ . Then since  $DG$  is parallel to  $AF$  and  $\angle EAF = \angle FAB$ ,  $DG$  will be parallel to  $AB$ .

$\therefore \triangle DEG$  and  $\triangle AFB$  are similar, and  $\angle EDG = \angle ABF = \frac{1}{3} \angle ABC$ . Similarly, it may be shown that  $\angle FDH$  is equal to  $\frac{1}{3} \angle ACB$ .

$\therefore \angle EDF = \frac{1}{3} (\angle ABC + \angle ACD + \angle BAC) = 60^\circ$ . In the same way,  $\angle DEF$  is shown to be equal to  $60^\circ$ .

$\therefore \triangle DEF$  is an equilateral triangle.

371. Proposed by W. S. HUGHES, Student, Williams College.

A right circular cone is cut by two parallel planes, one passing through the vertex, and each cutting both nappes. Are the straight lines which constitute the first section parallel to the asymptotes of the hyperbola forming the other section?

Solution by FRANK LOXLEY GRIFFIN, Ph. D., Assistant Professor of Mathematics, Williams College.

As the  $X$ -axis take the axis of the cone, and as the  $Y$ -axis the intersection of the first cutting plane with the plane through the vertex perpendicular to the  $X$ -axis. Then the equation of the cone is  $y^2 + z^2 = m^2 x^2$ , where  $m$  denotes the tangent of one-half the vertex angle of the cone. In the plane  $XOZ$  rotate the axes  $OX$  and  $OZ$  through an angle  $a$ , such that  $OX$  shall be in the first cutting plane. Then the cone and the cutting planes are given respectively by

$$(1) \quad y^2 + (x' \sin a + z' \cos a)^2 = m^2 (x' \cos a - z' \sin a)^2,$$

$$(2), (3) \quad z' = 0, \quad z' = d,$$

where  $d$  is the distance between the cutting planes. Now the straight lines of the first section are given by (2) and the equation obtained by making  $z' = 0$  in (1), say  $B^2 x'^2 - y^2 = 0$ . And the hyperbola is given by (3) and an